

### Laws


Multiple Use Sustained Yield Act – 1960


Forest and Rangeland Renewable
 Resources Planning Act – 1974

Forest Management Act - 1976

### Variability


#### North to South






### Variability

#### Lower elevation to higher elevation






### **Bailey's Sections and Subsections - Clearwater**




### Bailey's Sections and Subsections – Nez Perce



#### Idaho Batholith Breaklands

- Adjacent to rivers and larger streams
- Steep ground
- North and south aspects have different ecology



#### **Bitterroot Mountains Breaklands**

- Adjacent to rivers and larger streams
- Steep ground
- North and south aspects have different ecology





### **Uplands**

- Higher elevation
- Rolling topography

Deeper soils

Moderate climate




### **Idaho Batholith Subalpine**

Higher elevation

Colder climate

Mixed topography



### **Bitterroot Mountains Subalpine**

Higher elevation

Colder climate

Mixed topography



### **Forest Types**

**Examples:** 

PP – ponderosa pine

DF – Douglas-fir

**GF-** grand fir

WRC – western redcedar

These include mixes of these species, but they make up the character of the stand.







# Size Classes: Seral shrubs Trees:

0" to 5"
5" to 15"
15" to 20"
20" and greater








## Structure<br/>Single-storied

**Two-storied** 

**Multi-storied** 



# Dead Wood Both standing and down



Disturbance Regimes

- Fire
- Insects
- Disease
- ClimateVariation



#### **Patch Sizes**

Scale of disturbance – varies by subsection, driven by fire regimes and insect/disease cycles.

- 1. Breaklands large, every 5 to 50 years, low to moderate severity
- 2. Uplands small to large, every 50 to 200+ years, moderate to high severity
- 3. Subalpine small to large, every 50 to 150+ years, moderate to high severity

Historic Range of Variation shows what conditions provide for sustainable systems.

**Desired Conditions.** 

Current vegetation
Compared to HRV leads to
Objectives

### Bitterroot Forest Reserve 1898 Report by John Leiberg

- Identified 12 forest species
- The white pine type covered 30% of the Bitterroot Reserve – mostly on the Clearwater
- The ponderosa pine type covered 32% of the Reserve – mostly on the Nez Perce
- Larch was also a major component with white pine, and extended beyond the range of white pine
- 40% of the reserve had burned in the previous 50 years.


### **Changes In Forest Composition**

 Historically dominated by <u>potentially long-lived</u>, shade-intolerant, drought-tolerant, fire-adapted tree species:


white pine / larch / ponderosa pine / whitebark pine;

- <u>Currently</u> dominated by shade-tolerant, drought
   & fire sensitive, & <u>short-lived</u> late seral species:
  - grand fir / western hemlock / Douglas-fir / cedar


### Changes in Forest Cover Types – Clearwater NF



### Changes in Forest Cover Types – Nez Perce NF



## Historic and Current Age Class Distribution




### **Causes of Forest Change**

- White Pine Blister Rust (exotic disease);
- Big fires in early 20<sup>th</sup> century;
- Fire suppression;
- Past timber harvest practices

#### **Climate Variation**

 Climate goes through cycles. We are currently in a warm, dry phase, though in the 1940s through 1970s, we were in a cool, moist variation. Fire occurrence and extent have been tied to these climate variations.



Pacific Decadal Oscillation phase

From David L. Peterson, USDA Forest Service, PNW Research Station

### Implications of Forest Changes: So What ??

- Lower resilience to drought;
- Increasing risk from Insects & Diseases
  - Root pathogens / bark beetles / defoliators;
  - More insect & disease mortality = more rapid build up of dead fuels & ladder fuels;
- Increasing risk of severe wildfires;
- Less resilience to climate variations.

# Proposed Forest Vegetation Plan Components

- Desired Conditions
  - Species Resilient to Drought
  - Insect and Disease Resistant
  - Species that are Fire resistant
  - Better size class distribution
  - Less dense stands

# Proposed Forest Vegetation Plan Components

#### **Objectives**

 Treat about 250,000 acres across the Forest, on all settings, to facilitate moving the vegetation from current conditions toward desired conditions

#### **Guidelines**

 Favor long-lived, shade intolerant, drought tolerant, fire-adapted species

### Connecting 'Vegetation' with 'Habitat'

Food
Shelter
Reproduction


**Species Diversity** 




### **Seral Grass/Shrubs and Small Trees**



#### Dense Small Trees and Trees 15" to 20"



### Trees 15" to 20" and Over 20"



### Works-in-Progress

- Identification of Species of Conservation
   Concern
- Modeling and Mapping of Specific Habitats
- Motorized Influences

### **Forest-wide Plan Components**

#### **Habitat Needs Not Related to Vegetation**

- Security --- motorized disturbance
- Habitat access ---bat gates
- Limiting disease transmission --- bighorn sheep

### **Idaho Fish and Game**

#### **Role in Forest Plan Revision Process**



### Fish and Game Mission Statement

"All wildlife, including all wild animals, wild birds, and fish, within the state of Idaho, is hereby declared to be the property of the state of Idaho. It shall be preserved, protected, perpetuated, and managed. It shall be only captured or taken at such times or places, under such conditions, or by such means, or in such manner, as will preserve, protect, and perpetuate such wildlife, and provide for the citizens of this state and, as by law permitted to others, continued supplies of such wildlife for hunting, fishing and trapping."



### Fish and Game's Role

in Forest Plan Revision

Partnership . . .



### **Partnership Goals**

#### **My Roles**

- ID Team member
  - Inform the Assessment data, surveys, trends
  - Assist with interpretation of science into Plan
  - Meet F&G wildlife objectives, goals, mandates

Collaborative Member

# Non- Forest Vegetation Overview



### Non-Forest Components

- Shrublands
- Grasslands
- Rock outcroppings
- Other inclusions



Approx 15% of breaklands and <5% of uplands</li>

### **Non-Forest Uplands**

 Shrublands dominated by ninebark, ocean spray, alder, snowberry, menziesia and others species.



 Grasslands and dry meadows dominated by Idaho fescue, mountain brome, blue wildrye, sedges and other species



#### **Non-Forest Breaklands**

 Shrublands dominated by mountain mahogany, hackberry, and smooth sumac.

 Grasslands dominated by bluebunch wheatgrass, Idaho fescue, prairie junegrass, Sandburg's bluegrass and a variety of native forbs.





#### The Diverse Value of Grasslands



# Grasslands provide habitat for rare plant species



## Adverse impacts (stressors) can alter grasslands

• Livestock intensity/practices (primarily historic)

• "Un-natural" fire effects

Invasive weeds



### Current grassland conditions on the Forests

Limited data, representative areas indicate:

>50% high native integrity

<25% low native integrity</li>





#### **Invasive Weeds Pose a Serious Threat**

Warm, dry canyon environments are especially vulnerable to invasive weeds






- 425,000 acres infested
- 80 different species
- 500,000 acres weed free

## **Invasive Weeds Alter Ecological Processes**


- Displace native vegetation
- Wildlife habitat and forage
- Fire frequency & intensity
- Surface erosion





#### **Invasive Weeds of Highest Concern**

- Cheatgrass, Medusahead Rye, and Wiregrass
- Yellow/Orange Hawkweed
- Spotted Knapweed
- Rush Skeletonweed
- Yellow Starthistle
- Sulfur Cinquefoil



### Proposed Forest Plan Components Non-Forest Vegetation

- Desired Condition
  - Native plant communities
  - No new invasive weed species
- Objectives
  - Cooperation
  - Integrated Pest Management
  - Reduce conifer encroachment
  - Sensitive plant habitat protection and maintenance
- Guidelines:
  - Specific livestock grazing guidelines
  - Weed free feed and seed